

We see that the graph of $y = A \sin(\omega x - \phi) = A \sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$ is the same as the graph of $y = A \sin(\omega x)$, except that it has been shifted $\frac{\phi}{\omega}$ units (to the right if $\phi > 0$ and to the left if $\phi < 0$). This number $\frac{\phi}{\omega}$ is called the **phase shift** of the graph of $y = A \sin(\omega x - \phi)$.

For the graphs of $y = A \sin(\omega x - \phi)$ or $y = A \cos(\omega x - \phi)$, $\omega > 0$,

$$\text{Amplitude} = |A| \quad \text{Period} = T = \frac{2\pi}{\omega} \quad \text{Phase shift} = \frac{\phi}{\omega}$$

The phase shift is to the left if $\phi < 0$ and to the right if $\phi > 0$.

EXAMPLE 1

Finding the Amplitude, Period, and Phase Shift of a Sinusoidal Function and Graphing It

Find the amplitude, period, and phase shift of $y = 3 \sin(2x - \pi)$ and graph the function.

Solution Comparing

$$y = 3 \sin(2x - \pi) = 3 \sin\left[2\left(x - \frac{\pi}{2}\right)\right]$$

to

$$y = A \sin(\omega x - \phi) = A \sin\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$$

we find that $A = 3$, $\omega = 2$, and $\phi = \pi$. The graph is a sine curve with amplitude $|A| = 3$, period $T = \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi$, and phase shift $= \frac{\phi}{\omega} = \frac{\pi}{2}$.

The graph of $y = 3 \sin(2x - \pi)$ will lie between -3 and 3 on the y -axis. One cycle will begin at $x = \frac{\phi}{\omega} = \frac{\pi}{2}$ and end at $x = \frac{\phi}{\omega} + \frac{2\pi}{\omega} = \frac{\pi}{2} + \pi = \frac{3\pi}{2}$. To find

the five key points, we divide the interval $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ into four subintervals, each of length $\pi \div 4 = \frac{\pi}{4}$, by finding the following values of x :

$$\frac{\pi}{2} \quad \frac{\pi}{2} + \frac{\pi}{4} = \frac{3\pi}{4} \quad \frac{3\pi}{4} + \frac{\pi}{4} = \pi \quad \pi + \frac{\pi}{4} = \frac{5\pi}{4} \quad \frac{5\pi}{4} + \frac{\pi}{4} = \frac{3\pi}{2}$$

initial value

2nd value

3rd value

4th value


final value

Use these values of x to determine the five key points on the graph:

$$\left(\frac{\pi}{2}, 0\right), \left(\frac{3\pi}{4}, 3\right), (\pi, 0), \left(\frac{5\pi}{4}, -3\right), \left(\frac{3\pi}{2}, 0\right)$$


We plot these five points and fill in the graph of the sine function as shown in Figure 73(a). Extending the graph in each direction, we obtain Figure 73(b).

Figure 73



The graph of $y = 3 \sin(2x - \pi) = 3 \sin\left[2\left(x - \frac{\pi}{2}\right)\right]$ may also be obtained using transformations. See Figure 74.

Figure 74



To graph a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$, we first graph the function $y = A \sin(\omega x - \phi)$ and then apply a vertical shift.

EXAMPLE 2**Finding the Amplitude, Period, and Phase Shift of a Sinusoidal Function and Graphing It**

Find the amplitude, period, and phase shift of $y = 2 \cos(4x + 3\pi) + 1$ and graph the function.

Solution We begin by graphing $y = 2 \cos(4x + \pi)$. Comparing

$$y = 2 \cos(4x + 3\pi) = 2 \cos\left[4\left(x + \frac{3\pi}{4}\right)\right]$$

to

$$y = A \cos(\omega x - \phi) = A \cos\left[\omega\left(x - \frac{\phi}{\omega}\right)\right]$$

NOTE We can also find the interval defining one cycle by solving the inequality

$$0 \leq 4x + 3\pi \leq 2\pi$$

Then

$$\begin{aligned} -3\pi &\leq 4x \leq -\pi \\ -\frac{3\pi}{4} &\leq x \leq -\frac{\pi}{4} \end{aligned}$$

we see that $A = 2$, $\omega = 4$, and $\phi = -3\pi$. The graph is a cosine curve with amplitude $|A| = 2$, period $T = \frac{2\pi}{\omega} = \frac{2\pi}{4} = \frac{\pi}{2}$, and phase shift $= \frac{\phi}{\omega} = -\frac{3\pi}{4}$.

The graph of $y = 2 \cos(4x + 3\pi)$ will lie between -2 and 2 on the y -axis. One cycle will begin at $x = \frac{\phi}{\omega} = -\frac{3\pi}{4}$ and end at $x = \frac{\phi}{\omega} + \frac{2\pi}{\omega} = -\frac{3\pi}{4} + \frac{\pi}{2} = -\frac{\pi}{4}$.

To find the five key points, we divide the interval $\left[-\frac{3\pi}{4}, -\frac{\pi}{4}\right]$ into four subintervals, each of the length $\frac{\pi}{2} \div 4 = \frac{\pi}{8}$, by finding the following values.


$$\begin{array}{cccccc} -\frac{3\pi}{4} & -\frac{3\pi}{4} + \frac{\pi}{8} = -\frac{5\pi}{8} & -\frac{5\pi}{8} + \frac{\pi}{8} = -\frac{\pi}{2} & -\frac{\pi}{2} + \frac{\pi}{8} = -\frac{3\pi}{8} & -\frac{3\pi}{8} + \frac{\pi}{8} = -\frac{\pi}{4} \\ \text{initial value} & \text{2nd value} & \text{3rd value} & \text{4th value} & \text{final value} \end{array}$$

The five key points on the graph of $y = 2 \cos(4x + \pi)$ are

$$\left(-\frac{3\pi}{4}, 2\right), \left(-\frac{5\pi}{8}, 0\right), \left(-\frac{\pi}{2}, -2\right), \left(-\frac{3\pi}{8}, 0\right), \left(-\frac{\pi}{4}, 2\right)$$


We plot these five points and fill in the graph of the cosine function as shown in Figure 75(a). Extending the graph in each direction, we obtain Figure 75(b), the graph of $y = 2 \cos(4x + \pi)$. A vertical shift up 1 unit gives the final graph. See Figure 75(c).

Figure 75



The graph of $y = 2 \cos(4x + 3\pi) + 1 = 2 \cos\left[4\left(x + \frac{3\pi}{4}\right)\right] + 1$ may also be obtained using transformations. See Figure 76.

Figure 76



Now Work PROBLEM 3

SUMMARY Steps for Graphing Sinusoidal Functions $y = A \sin(\omega x - \phi) + B$ or $y = A \cos(\omega x - \phi) + B$

STEP 1: Determine the amplitude $|A|$ and period $T = \frac{2\pi}{\omega}$.

STEP 2: Determine the starting point of one cycle of the graph, $\frac{\phi}{\omega}$.

STEP 3: Determine the ending point of one cycle of the graph, $\frac{\phi}{\omega} + \frac{2\pi}{\omega}$.

STEP 4: Divide the interval $\left[\frac{\phi}{\omega}, \frac{\phi}{\omega} + \frac{2\pi}{\omega}\right]$ into four subintervals, each of length $\frac{2\pi}{\omega} \div 4$.

STEP 5: Use the endpoints of the subintervals to find the five key points on the graph.

STEP 6: Fill in one cycle of the graph.

STEP 7: Extend the graph in each direction to make it complete.

STEP 8: If $B \neq 0$, apply a vertical shift.


2 Find a Sinusoidal Function from Data

A review of scatter diagrams from Appendix A, Section A.8, may be helpful.

Scatter diagrams of data sometimes take the form of a sinusoidal function. Let's look at an example.

The data given in Table 11 represent the average monthly temperatures in Denver, Colorado. Since the data represent *average* monthly temperatures collected over many years, the data will not vary much from year to year and so will essentially repeat each year. In other words, the data are periodic. Figure 77 shows the scatter diagram of these data repeated over 2 years, where $x = 1$ represents January, $x = 2$ represents February, and so on.


Table 11



Month, x	Average Monthly Temperature, °F
January, 1	29.7
February, 2	33.4
March, 3	39.0
April, 4	48.2
May, 5	57.2
June, 6	66.9
July, 7	73.5
August, 8	71.4
September, 9	62.3
October, 10	51.4
November, 11	39.0
December, 12	31.0

SOURCE: U.S. National Oceanic and Atmospheric Administration

Figure 77



Notice that the scatter diagram looks like the graph of a sinusoidal function. We choose to fit the data to a sine function of the form

$$y = A \sin(\omega x - \phi) + B$$

where A , B , ω , and ϕ are constants.

EXAMPLE 3

Finding a Sinusoidal Function from Temperature Data

Fit a sine function to the data in Table 11.

Solution

We begin with a scatter diagram of the data for one year. See Figure 78. The data will be fitted to a sine function of the form

$$y = A \sin(\omega x - \phi) + B$$

STEP 1: To find the amplitude A , we compute

$$\begin{aligned} \text{Amplitude} &= \frac{\text{largest data value} - \text{smallest data value}}{2} \\ &= \frac{73.5 - 29.7}{2} = 21.9 \end{aligned}$$

To see the remaining steps in this process, we superimpose the graph of the function $y = 21.9 \sin x$, where x represents months, on the scatter diagram.

Figure 78




Figure 79




Figure 79 shows the two graphs. To fit the data, the graph needs to be shifted vertically, shifted horizontally, and stretched horizontally.

STEP 2: We determine the vertical shift by finding the average of the highest and lowest data values.

$$\text{Vertical shift} = \frac{73.5 + 29.7}{2} = 51.6$$

Now we superimpose the graph of $y = 21.9 \sin x + 51.6$ on the scatter diagram. See Figure 80.

We see that the graph needs to be shifted horizontally and stretched horizontally.

STEP 3: It is easier to find the horizontal stretch factor first. Since the temperatures repeat every 12 months, the period of the function is $T = 12$. Since

$$T = \frac{2\pi}{\omega} = 12, \text{ we find}$$

$$\omega = \frac{2\pi}{12} = \frac{\pi}{6}$$

Now we superimpose the graph of $y = 21.9 \sin\left(\frac{\pi}{6}x\right) + 51.6$ on the scatter diagram. See Figure 81. We see that the graph still needs to be shifted horizontally.

Figure 80





Figure 81



STEP 4: To determine the horizontal shift, we use the period $T = 12$ and divide the interval $[0, 12]$ into four subintervals of length $12 \div 4 = 3$:

$$[0, 3], [3, 6], [6, 9], [9, 12]$$

The sine curve is increasing on the interval $(0, 3)$ and is decreasing on the interval $(3, 9)$, so a local maximum occurs at $x = 3$. The data indicate that a maximum occurs at $x = 7$ (corresponding to July's temperature), so we must shift the graph of the function 4 units to the right by replacing x by $x - 4$. Doing this, we obtain


$$y = 21.9 \sin\left(\frac{\pi}{6}(x - 4)\right) + 51.6$$

Multiplying out, we find that a sine function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data is

$$y = 21.9 \sin\left(\frac{\pi}{6}x - \frac{2\pi}{3}\right) + 51.6$$

The graph of $y = 21.9 \sin\left(\frac{\pi}{6}x - \frac{2\pi}{3}\right) + 51.6$ and the scatter diagram of the data are shown in Figure 82.

Figure 82



The steps to fit a sine function

$$y = A \sin(\omega x - \phi) + B$$

to sinusoidal data follow:

Steps for Fitting Data to a Sine Function $y = A \sin(\omega x - \phi) + B$

STEP 1: Determine A , the amplitude of the function.

$$\text{Amplitude} = \frac{\text{largest data value} - \text{smallest data value}}{2}$$

STEP 2: Determine B , the vertical shift of the function.

$$\text{Vertical shift} = \frac{\text{largest data value} + \text{smallest data value}}{2}$$

STEP 3: Determine ω . Since the period T , the time it takes for the data to repeat, is $T = \frac{2\pi}{\omega}$, we have

$$\omega = \frac{2\pi}{T}$$

STEP 4: Determine the horizontal shift of the function by using the period of the data. Divide the period into four subintervals of equal length. Determine the x -coordinate for the maximum of the sine function and the x -coordinate for the maximum value of the data. Use this

information to determine the value of the phase shift, $\frac{\phi}{\omega}$.

Now Work PROBLEM 29(a)–(c)


Let's look at another example. Since the number of hours of sunlight in a day cycles annually, the number of hours of sunlight in a day for a given location can be modeled by a sinusoidal function.

The longest day of the year (in terms of hours of sunlight) occurs on the day of the summer solstice. For locations in the northern hemisphere, the summer solstice is the time when the sun is farthest north. In 2005, the summer solstice occurred on June 21 (the 172nd day of the year) at 2:46 AM EDT. The shortest day of the year occurs on the day of the winter solstice. The winter solstice is the time when the Sun is farthest south (again, for locations in the northern hemisphere). In 2005, the winter solstice occurred on December 21 (the 355th day of the year) at 1:35 PM (EST).

EXAMPLE 4

Finding a Sinusoidal Function for Hours of Daylight

According to the *Old Farmer's Almanac*, the number of hours of sunlight in Boston on the summer solstice is 15.30 and the number of hours of sunlight on the winter solstice is 9.08.

- Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
-  Draw a graph of the function found in part (a).
- Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac* and compare it to the results found in part (b).

Source: The *Old Farmer's Almanac*, www.almanac.com/rise

Solution

$$\begin{aligned} \text{(a) STEP 1: Amplitude} &= \frac{\text{largest data value} - \text{smallest data value}}{2} \\ &= \frac{15.30 - 9.08}{2} = 3.11 \end{aligned}$$

$$\begin{aligned}\text{STEP 2: Vertical shift} &= \frac{\text{largest data value} + \text{smallest data value}}{2} \\ &= \frac{15.30 + 9.08}{2} = 12.19\end{aligned}$$

STEP 3: The data repeat every 365 days. Since $T = \frac{2\pi}{\omega} = 365$, we find

$$\omega = \frac{2\pi}{365}$$

So far, we have $y = 3.11 \sin\left(\frac{2\pi}{365}x - \phi\right) + 12.19$.

STEP 4: To determine the horizontal shift, we use the period $T = 365$ and divide the interval $[0, 365]$ into four subintervals of length $365 \div 4 = 91.25$:

$$[0, 91.25], [91.25, 182.5], [182.5, 273.75], [273.75, 365]$$

The sine curve is increasing on the interval $(0, 91.25)$ and is decreasing on the interval $(91.25, 273.75)$, so a local maximum occurs at $x = 91.25$. Since the maximum occurs on the summer solstice at $x = 172$, we must shift the graph of the function $172 - 91.25 = 80.75$ units to the right by replacing x by $x - 80.75$. Doing this, we obtain

$$y = 3.11 \sin\left(\frac{2\pi}{365}(x - 80.75)\right) + 12.19$$

Multiplying out, we find that a sine function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data is

$$y = 3.11 \sin\left(\frac{2\pi}{365}x - \frac{323\pi}{730}\right) + 12.19$$

(b) To predict the number of hours of daylight on April 1, we let $x = 91$ in the function found in part (a) and obtain

$$\begin{aligned}y &= 3.11 \sin\left(\frac{2\pi}{365} \cdot 91 - \frac{323\pi}{730}\right) + 12.19 \\ &\approx 12.74\end{aligned}$$


So we predict that there will be about 12.74 hours = 12 hours, 44 minutes of sunlight on April 1 in Boston.



(c) The graph of the function found in part (a) is given in Figure 83.

(d) According to the *Old Farmer's Almanac*, there will be 12 hours 45 minutes of sunlight on April 1 in Boston.

Figure 83



Now Work PROBLEM 35

Certain graphing utilities (such as a TI-83, TI-84 Plus, and TI-86) have the capability of finding the sine function of best fit for sinusoidal data. At least four data points are required for this process.

EXAMPLE 5

Finding the Sine Function of Best Fit

Use a graphing utility to find the sine function of best fit for the data in Table 15. Graph this function with the scatter diagram of the data.

Solution

Enter the data from Table 15 and execute the SINE REGression program. The result is shown in Figure 84.

The output that the utility provides shows the equation

$$y = a \sin(bx + c) + d$$

The sinusoidal function of best fit is

$$y = 21.15 \sin(0.55x - 2.35) + 51.19$$

where x represents the month and y represents the average temperature.

Figure 85 shows the graph of the sinusoidal function of best fit on the scatter diagram.

Figure 84





Figure 85



Now Work PROBLEMS 29(d) AND (e)

2.6 Assess Your Understanding

Concepts and Vocabulary

- For the graph of $y = A \sin(\omega x - \phi)$, the number $\frac{\phi}{\omega}$ is called the _____.
- True or False** Only two data points are required by a graphing utility to find the sine function of best fit.

Skill Building

In Problems 3–14, find the amplitude, period, and phase shift of each function. Graph each function. Be sure to label key points. Show at least two periods.

- $y = 4 \sin(2x - \pi)$
- $y = 3 \sin(3x - \pi)$
- $y = 2 \cos\left(3x + \frac{\pi}{2}\right)$
- $y = 3 \cos(2x + \pi)$
- $y = -3 \sin\left(2x + \frac{\pi}{2}\right)$
- $y = -2 \cos\left(2x - \frac{\pi}{2}\right)$
- $y = 4 \sin(\pi x + 2) - 5$
- $y = 2 \cos(2\pi x + 4) + 4$
- $y = 3 \cos(\pi x - 2) + 5$
- $y = 2 \cos(2\pi x - 4) - 1$
- $y = -3 \sin\left(-2x + \frac{\pi}{2}\right)$
- $y = -3 \cos\left(-2x + \frac{\pi}{2}\right)$

In Problems 15–18, write the equation of a sine function that has the given characteristics.

- | | | | |
|----------------------------|-------------------------|-----------------------------|-------------------|
| 15. Amplitude: 2 | 16. Amplitude: 3 | 17. Amplitude: 3 | 18. Amplitude: 2 |
| Period: π | Period: $\frac{\pi}{2}$ | Period: 3π | Period: π |
| Phase shift: $\frac{1}{2}$ | Phase shift: 2 | Phase shift: $-\frac{1}{3}$ | Phase shift: -2 |

Applications and Extensions

In Problems 19–26, apply the methods of this and the previous section to graph each function. Be sure to label key points and show at least two periods.

- $y = 2 \tan(4x - \pi)$
- $y = \frac{1}{2} \cot(2x - \pi)$
- $y = 3 \csc\left(2x - \frac{\pi}{4}\right)$
- $y = \frac{1}{2} \sec(3x - \pi)$
- $y = -\cot\left(2x + \frac{\pi}{2}\right)$
- $y = -\tan\left(3x + \frac{\pi}{2}\right)$
- $y = -\sec(2\pi x + \pi)$
- $y = -\csc\left(-\frac{1}{2}\pi x + \frac{\pi}{4}\right)$

- 27. Alternating Current (ac) Circuits** The current I , in amperes, flowing through an ac (alternating current) circuit at time t , in seconds, is

$$I(t) = 120 \sin\left(30\pi t - \frac{\pi}{3}\right), \quad t \geq 0$$


What is the period? What is the amplitude? What is the phase shift? Graph this function over two periods.

- 28. Alternating Current (ac) Circuits** The current I , in amperes, flowing through an ac (alternating current) circuit at time t , in seconds, is

$$I(t) = 220 \sin\left(60\pi t - \frac{\pi}{6}\right), \quad t \geq 0$$


What is the period? What is the amplitude? What is the phase shift? Graph this function over two periods.

- 29. Monthly Temperature** The following data represent the average monthly temperatures for Juneau, Alaska.



Month, x	Average Monthly Temperature, °F
January, 1	24.2
February, 2	28.4
March, 3	32.7
April, 4	39.7
May, 5	47.0
June, 6	53.0
July, 7	56.0
August, 8	55.0
September, 9	49.4
October, 10	42.2
November, 11	32.0
December, 12	27.1


SOURCE: U.S. National Oceanic and Atmospheric Administration



Month, x	Average Monthly Temperature, °F
January, 1	34.6
February, 2	37.5
March, 3	47.2
April, 4	56.5
May, 5	66.4
June, 6	75.6
July, 7	80.0
August, 8	78.5
September, 9	71.3
October, 10	59.7
November, 11	49.8
December, 12	39.4

SOURCE: U.S. National Oceanic and Atmospheric Administration


- 31. Monthly Temperature** The following data represent the average monthly temperatures for Indianapolis, Indiana.



Month, x	Average Monthly Temperature, °F
January, 1	25.5
February, 2	29.6
March, 3	41.4
April, 4	52.4
May, 5	62.8
June, 6	71.9
July, 7	75.4
August, 8	73.2
September, 9	66.6
October, 10	54.7
November, 11	43.0
December, 12	30.9


SOURCE: U.S. National Oceanic and Atmospheric Administration

- (a) Draw a scatter diagram of the data for one period.
 (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
 (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
 (d) Use a graphing utility to find the sinusoidal function of best fit.
 (e) Draw the sinusoidal function of best fit on a scatter diagram of the data.
- 30. Monthly Temperature** The following data represent the average monthly temperatures for Washington, D.C.
- (a) Draw a scatter diagram of the data for one period.
 (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
 (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
 (d) Use a graphing utility to find the sinusoidal function of best fit.
 (e) Graph the sinusoidal function of best fit on a scatter diagram of the data.
- (a) Draw a scatter diagram of the data for one period.
 (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
 (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
 (d) Use a graphing utility to find the sinusoidal function of best fit.
 (e) Graph the sinusoidal function of best fit on a scatter diagram of the data.
- 32. Monthly Temperature** The data on the following page represent the average monthly temperatures for Baltimore, Maryland.
- (a) Draw a scatter diagram of the data for one period.




Month, x	Average Monthly Temperature, °F
January, 1	31.8
February, 2	34.8
March, 3	44.1
April, 4	53.4
May, 5	63.4
June, 6	72.5
July, 7	77.0
August, 8	75.6
September, 9	68.5
October, 10	56.6
November, 11	46.8
December, 12	36.7

SOURCE: U.S. National Oceanic and Atmospheric Administration

- (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (c) Draw the sinusoidal function found in part (b) on the scatter diagram.
-  (d) Use a graphing utility to find the sinusoidal function of best fit.
- (e) Graph the sinusoidal function of best fit on a scatter diagram of the data.
- 33. Tides** Suppose that the length of time between consecutive high tides is approximately 12.5 hours. According to the National Oceanic and Atmospheric Administration, on Saturday, August 7, 2004, in Savannah, Georgia, high tide occurred at 3:38 AM (3.6333 hours) and low tide occurred at 10:08 AM (10.1333 hours). Water heights are measured as the amounts above or below the mean lower low water. The height of the water at high tide was 8.2 feet, and the height of the water at low tide was -0.6 foot.
- (a) Approximately when will the next high tide occur?
- (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (c) Draw a graph of the function found in part (b).
- (d) Use the function found in part (b) to predict the height of the water at the next high tide.
- 34. Tides** Suppose that the length of time between consecutive high tides is approximately 12.5 hours. According to the National Oceanic and Atmospheric Administration, on Saturday, August 7, 2004, in Juneau, Alaska, high tide occurred at 8:11 AM (8.1833 hours) and low tide occurred at 2:14 PM (14.2333 hours). Water heights are measured as the amounts above or below the mean lower low water. The height of the water at high tide was 13.2 feet, and the height of the water at low tide was 2.2 feet.
- (a) Approximately when will the next high tide occur?
- (b) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.

- (c) Draw a graph of the function found in part (b).
- (d) Use the function found in part (b) to predict the height of the water at the next high tide.

-  **35. Hours of Daylight** According to the *Old Farmer's Almanac*, in Miami, Florida, the number of hours of sunlight on the summer solstice of 2005 is 13.75 and the number of hours of sunlight on the winter solstice is 10.53.
- (a) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
- 36. Hours of Daylight** According to the *Old Farmer's Almanac*, in Detroit, Michigan, the number of hours of sunlight on the summer solstice of 2005 is 15.30 and the number of hours of sunlight on the winter solstice is 9.08.
- (a) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
- 37. Hours of Daylight** According to the *Old Farmer's Almanac*, in Anchorage, Alaska, the number of hours of sunlight on the summer solstice of 2005 is 19.42 and the number of hours of sunlight on the winter solstice is 5.47.
- (a) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).
- 38. Hours of Daylight** According to the *Old Farmer's Almanac*, in Honolulu, Hawaii, the number of hours of sunlight on the summer solstice of 2005 is 13.43 and the number of hours of sunlight on the winter solstice is 10.85.
- (a) Find a sinusoidal function of the form $y = A \sin(\omega x - \phi) + B$ that fits the data.
- (b) Use the function found in part (a) to predict the number of hours of sunlight on April 1, the 91st day of the year.
- (c) Draw a graph of the function found in part (a).
- (d) Look up the number of hours of sunlight for April 1 in the *Old Farmer's Almanac*, and compare the actual hours of daylight to the results found in part (c).

Discussion and Writing

- 39.** Explain how the amplitude and period of a sinusoidal graph are used to establish the scale on each coordinate axis.
- 40.** Find an application in your major field that leads to a sinusoidal graph. Write a paper about your findings.